

# Energy loss in Unstable Quark-Gluon Plasma



**Stanisław Mrówczyński**

*Jan Kochanowski University, Kielce, Poland  
& National Centre of Nuclear Research, Warsaw, Poland*

in collaboration with



**Margaret Carrington & Katarzyna Deja**

# Motivation - jet quenching



Away-side jet is suppressed  
in central collisions


# Scenario of relativistic heavy-ion collisions



before

QGP is out of equilibrium at the collision early stage

# Anisotropic QGP



Anisotropic QGP is unstable due to magnetic plasma modes

## Energy loss in unstable QGP

- ▶ Is  $dE/dx$  in unstable QGP sizeable?  
Yes!
- ▶ How to compute  $dE/dx$  in unstable QGP?  
Solve initial value problem!

# A test parton in QGP

**Wong's equation of motion** (Hard Loop Approximation)

$$\begin{cases} \frac{dx^\mu(\tau)}{d\tau} = u^\mu(\tau) \\ \frac{dp^\mu(\tau)}{d\tau} = gQ_a(\tau) F_a^{\mu\nu}(x(\tau)) u_\nu(\tau) \\ \frac{dQ_a(\tau)}{d\tau} = -gf^{abc} p_\mu(\tau) A_b^\mu(x(\tau)) Q_c(\tau) \end{cases}$$

## Simplifications

Gauge condition:  $p_\mu(\tau) A_b^\mu(x(\tau)) = 0 \Rightarrow Q_a(\tau) = \text{const}$

Parton travels with constant velocity:  $u^\mu = (\gamma, \gamma \mathbf{v}) = \text{const}$

## Parton's energy loss

$$\frac{dE(t)}{dt} = gQ_a \mathbf{E}_a(t, \mathbf{r}(t)) \cdot \mathbf{v}$$

induced & spontaneously  
generated chromoelectric field

parton's current:  $\mathbf{j}_a(t, \mathbf{r}) = gQ_a \mathbf{v} \delta^{(3)}(\mathbf{r} - \mathbf{v}t)$


$$\frac{dE(t)}{dt} = \int d^3r \mathbf{E}_a(t, \mathbf{r}) \cdot \mathbf{j}_a(t, \mathbf{r})$$

# Initial value problem

One-sided Fourier transformation

$$\left\{ \begin{array}{l} f(\omega, \mathbf{k}) = \int_0^{\infty} dt \int d^3 r e^{i(\omega t - \mathbf{k} \cdot \mathbf{r})} f(t, \mathbf{r}) \\ f(t, \mathbf{r}) = \int_{-\infty+i\sigma}^{\infty+i\sigma} \frac{d\omega}{2\pi} \int \frac{d^3 k}{(2\pi)^3} e^{-i(\omega t - \mathbf{k} \cdot \mathbf{r})} f(\omega, \mathbf{k}) \end{array} \right.$$

$0 < \sigma \in R$



$$\mathbf{j}_a(t, \mathbf{r}) = gQ_a \mathbf{v} \delta^{(3)}(\mathbf{r} - \mathbf{v}t) \Rightarrow \mathbf{j}_a(\omega, \mathbf{k}) = \frac{igQ_a \mathbf{v}}{\omega - \mathbf{k} \cdot \mathbf{v}}$$

$$\frac{dE(t)}{dt} = gQ_a \int_{-\infty+i\sigma}^{\infty+i\sigma} \frac{d\omega}{2\pi} \int \frac{d^3 k}{(2\pi)^3} e^{-i(\omega - \mathbf{k} \cdot \mathbf{v})t} \mathbf{E}_a(\omega, \mathbf{k}) \cdot \mathbf{v}$$

# Induced Electric Field

Linearized Yang-Mills (Maxwell) equations (Hard Loop Approximation)

$$\begin{aligned} i\mathbf{k} \cdot \mathbf{D}(\omega, \mathbf{k}) &= \rho(\omega, \mathbf{k}), & i\mathbf{k} \cdot \mathbf{B}(\omega, \mathbf{k}) &= 0, \\ i\mathbf{k} \times \mathbf{E}(\omega, \mathbf{k}) &= i\omega \mathbf{B}(\omega, \mathbf{k}) + \mathbf{B}_0(\mathbf{k}), \\ i\mathbf{k} \times \mathbf{B}(\omega, \mathbf{k}) &= \mathbf{j}(\omega, \mathbf{k}) - i\omega \mathbf{E}(\omega, \mathbf{k}) - \mathbf{D}_0(\mathbf{k}) \end{aligned}$$

$$D^i(\omega, \mathbf{k}) = \epsilon^{ij}(\omega, \mathbf{k}) E^j(\omega, \mathbf{k})$$

Chromodielectric tensor

$$\epsilon^{ij}(\omega, \mathbf{k}) = \delta^{ij} + \frac{g^2}{2\omega} \int \frac{d^3 p}{(2\pi)^3} \frac{v^i}{\omega - \mathbf{k}\mathbf{v} + i0^+} \frac{\partial f(\mathbf{p})}{\partial p^l} \left[ \left(1 - \frac{\mathbf{k}\mathbf{v}}{\omega}\right) \delta^{lj} + \frac{k^l v^j}{\omega} \right] \quad \text{dynamical information}$$

$$E^i(\omega, \mathbf{k}) = -i(\Sigma^{-1})^{ij}(\omega, \mathbf{k}) [\omega \mathbf{j}(\omega, \mathbf{k}) + \mathbf{k} \times \mathbf{B}_0(\mathbf{k}) - \omega \mathbf{D}_0(\mathbf{k})]^j$$

$$\Sigma^{ij}(\omega, \mathbf{k}) \equiv -\mathbf{k}^2 \delta^{ij} + k^i k^j + \omega^2 \epsilon^{ij}(\omega, \mathbf{k})$$

## Energy-Loss formula

$$\frac{dE(t)}{dt} = gQ_a v^i \int_{-\infty+i\sigma}^{\infty+i\sigma} \frac{d\omega}{2\pi i} \int \frac{d^3 k}{(2\pi)^3} e^{-i(\omega - \bar{\omega})t} \\ \times (\Sigma^{-1})^{ij}(\omega, \mathbf{k}) \left[ \frac{igQ_a \omega \mathbf{v}}{\omega - \bar{\omega}} + \mathbf{k} \times \mathbf{B}_0(\mathbf{k}) - \omega \mathbf{D}_0(\mathbf{k}) \right]^j$$

$$\bar{\omega} \equiv \mathbf{k} \cdot \mathbf{v}$$

$$\Sigma^{ij}(\omega, \mathbf{k}) \equiv -\mathbf{k}^2 \delta^{ij} + k^i k^j + \omega^2 \epsilon^{ij}(\omega, \mathbf{k})$$

### Dispersion equation

$$\det[\Sigma(\omega, \mathbf{k})] = 0$$

## Initial values of the fields

Maxwell equations &  $\mathbf{j}_a(t, \mathbf{r}) = gQ_a \mathbf{v} \delta^{(3)}(\mathbf{r} - \mathbf{v}t)$



Initial values:

$$D_0^i(\mathbf{k}) = -igQ_a \bar{\omega} \epsilon^{ij}(\bar{\omega}, \mathbf{k}) (\Sigma^{-1})^{jk}(\bar{\omega}, \mathbf{k}) v^k$$

$$B_0^i(\mathbf{k}) = -igQ_a \epsilon^{ijk} k^j (\Sigma^{-1})^{kl}(\bar{\omega}, \mathbf{k}) v^l$$

When the test parton enters the plasma at  $t = 0$ , the instabilities are initiated.

## Energy-Loss formula

$$\begin{aligned}
 \frac{dE(t)}{dt} = & ig^2 C_R v^i v^l \int_{-\infty+i\sigma}^{\infty+i\sigma} \frac{d\omega}{2\pi i} \int \frac{d^3 k}{(2\pi)^3} e^{-i(\omega - \bar{\omega})t} (\Sigma^{-1})^{ij}(\omega, \mathbf{k}) \\
 & \times \left[ \underbrace{\frac{\omega \delta^{jl}}{\omega - \bar{\omega}}}_{\mathbf{j}(\omega, \mathbf{k})} - (k^j k^k - \mathbf{k}^2 \delta^{jk}) (\Sigma^{-1})^{kl}(\bar{\omega}, \mathbf{k}) + \omega \bar{\omega} \epsilon^{jk}(\bar{\omega}, \mathbf{k}) (\Sigma^{-1})^{kl}(\bar{\omega}, \mathbf{k}) \right]
 \end{aligned}$$

Averaging over parton's colors:  $\int dQ Q_a Q_b = C_2 \delta^{ab}$ ,  $C_2 \equiv \begin{cases} \frac{1}{2} & \text{for quark} \\ N_c & \text{for gluon} \end{cases}$

$$C_R = \begin{cases} C_2 \frac{N_c^2 - 1}{N_c} = \frac{N_c^2 - 1}{2N_c} & \text{for quark (R = F)} \\ C_2 = N_c & \text{for gluon (R = G)} \end{cases}$$

## Stable isotropic plasma

►  $\epsilon^{ij}(\omega, \mathbf{k}) = \epsilon_L(\omega, \mathbf{k}) \frac{k^i k^j}{\mathbf{k}^2} + \epsilon_T(\omega, \mathbf{k}) \left( \delta^{ij} - \frac{k^i k^j}{\mathbf{k}^2} \right)$

►  $(\Sigma^{-1})^{ij}(\omega, \mathbf{k}) = \frac{1}{\omega^2 \epsilon_L(\omega, \mathbf{k})} \frac{k^i k^j}{\mathbf{k}^2} + \frac{1}{\omega^2 \epsilon_T(\omega, \mathbf{k}) - \mathbf{k}^2} \left( \delta^{ij} - \frac{k^i k^j}{\mathbf{k}^2} \right)$

$$\begin{aligned} \frac{dE(t)}{dt} &= ig^2 C_R v^i v^l \int_{-\infty+i\sigma}^{\infty+i\sigma} \frac{d\omega}{2\pi i} \int \frac{d^3 k}{(2\pi)^3} e^{-i(\omega - \bar{\omega})t} (\Sigma^{-1})^{ij}(\omega, \mathbf{k}) \\ &\times \left[ \frac{\omega \delta^{jl}}{\omega - \bar{\omega}} - (k^j k^k - \mathbf{k}^2 \delta^{jk}) (\Sigma^{-1})^{kl}(\bar{\omega}, \mathbf{k}) + \omega \bar{\omega} \epsilon^{jk}(\bar{\omega}, \mathbf{k}) (\Sigma^{-1})^{kl}(\bar{\omega}, \mathbf{k}) \right] \end{aligned}$$


The only stationary contribution:  $\omega = \bar{\omega} \equiv \mathbf{k} \cdot \mathbf{v}$

$$\frac{dE(t)}{dt} = ig^2 C_R \int \frac{d^3 k}{(2\pi)^3} \frac{\bar{\omega}}{\mathbf{k}^2} \left[ \frac{1}{\epsilon_L(\bar{\omega}, \mathbf{k})} + \frac{\mathbf{k}^2 \mathbf{v}^2 - \bar{\omega}^2}{\bar{\omega}^2 \epsilon_T(\bar{\omega}, \mathbf{k}) - \mathbf{k}^2} \right]$$


equivalent to the standard result by Braaten & Thoma 13

# Energy loss in equilibrium QGP

$$\frac{dE(t)}{dt} = ig^2 C_R \int \frac{d^3 k}{(2\pi)^3} \frac{\bar{\omega}}{\mathbf{k}^2} \left[ \frac{1}{\varepsilon_L(\bar{\omega}, \mathbf{k})} + \frac{\mathbf{k}^2 \mathbf{v}^2 - \bar{\omega}^2}{\bar{\omega}^2 \varepsilon_T(\bar{\omega}, \mathbf{k}) - \mathbf{k}^2} \right]$$



## Extremely prolate system



$$f(\mathbf{p}) \sim \delta(p_x) \delta(p_y)$$

Collective modes

$$\det[\Sigma^{ij}(\omega, \mathbf{k})] = 0$$

$$\Sigma^{ij}(\omega, \mathbf{k}) \equiv -\mathbf{k}^2 \delta^{ij} + k^i k^j + \omega^2 \varepsilon^{ij}(\omega, \mathbf{k})$$

$$\varepsilon^{ij}(\omega, \mathbf{k}) = \delta^{ij} + \frac{g^2}{2\omega} \int \frac{d^3 p}{(2\pi)^3} \frac{v^i}{\omega - \mathbf{k}\mathbf{v} + i0^+} \frac{\partial f(\mathbf{p})}{\partial p^l} \left[ \left( 1 - \frac{\mathbf{k}\mathbf{v}}{\omega} \right) \delta^{ij} + \frac{k^l v^j}{\omega} \right]$$

Spectrum of collective modes

$$\left\{ \begin{array}{ll} \omega_1(\mathbf{k}) = \mu^2 + \mathbf{k}^2 & \mathbf{n} \equiv (0,0,1) \\ \omega_2(\mathbf{k}) = \mu^2 + (\mathbf{k} \cdot \mathbf{n})^2 & \\ \omega_{\pm}(\mathbf{k}) = \frac{1}{2} \left( \mathbf{k}^2 + (\mathbf{k} \cdot \mathbf{n})^2 \pm \sqrt{\mathbf{k}^4 + (\mathbf{k} \cdot \mathbf{n})^4 + 4\mu^2 \mathbf{k}^2 - 4\mu^2 (\mathbf{k} \cdot \mathbf{n})^2 - 2\mathbf{k}^2 (\mathbf{k} \cdot \mathbf{n})^2} \right) & \end{array} \right.$$

# Unstable chromomagnetic mode

stationary state

$$A(t) = A_0 + \delta A(t)$$



fluctuation

Instability

$$\delta A(t) \propto e^{\text{Im } \omega t}$$

$$-\frac{\omega_-^2(\mathbf{k})}{\mu^2}$$

$$\text{Im } \omega > 0$$



$$\frac{dE(t)}{dt} \sim \int \frac{d^3 k}{(2\pi)^3} e^{\text{Im } \omega t} \dots$$

## Energy loss in extremely prolate system

$$\frac{dE(t)}{dt} = ig^2 C_R v^i v^l \int_{-\infty+i\sigma}^{\infty+i\sigma} \frac{d\omega}{2\pi i} \int \frac{d^3 k}{(2\pi)^3} e^{-i(\omega - \bar{\omega})t} (\Sigma^{-1})^{ij}(\omega, \mathbf{k}) \\ \times \left[ \frac{\omega \delta^{jl}}{\omega - \bar{\omega}} - (k^j k^k - \mathbf{k}^2 \delta^{jk}) (\Sigma^{-1})^{kl}(\bar{\omega}, \mathbf{k}) + \omega \bar{\omega} \epsilon^{jk}(\bar{\omega}, \mathbf{k}) (\Sigma^{-1})^{kl}(\bar{\omega}, \mathbf{k}) \right]$$

► Inversion of matrix  $\Sigma$  depending on  $\mathbf{k}$  and  $\mathbf{n}$

$$\Sigma = aA + bB + cC + dD \quad \Sigma^{-1} = \alpha A + \beta B + \gamma C + \delta D$$



basis of matrices

$$\left\{ \begin{array}{l} A^{ij} = \delta^{ij} - \frac{k^i k^j}{\mathbf{k}^2}, \quad B^{ij} = \frac{k^i k^j}{\mathbf{k}^2}, \\ C^{ij} = \frac{n_T^i n_T^j}{\mathbf{n}_T^2}, \quad D^{ij} = n_T^i k^j + k^i n_T^j \end{array} \right. \quad n_T^i \equiv \left( \delta^{ij} - \frac{k^i k^j}{\mathbf{k}^2} \right) n^j$$

$$\Sigma \Sigma^{-1} = 1 \quad \Rightarrow \quad \alpha, \beta, \gamma, \delta$$

## Energy loss in extremely prolate system cont.

$$\frac{1}{g^2 \mu^2} \frac{dE}{dt}$$



Debye mass

$$\mu^2 \equiv g^2 \int \frac{d^3 p}{(2\pi)^3} \frac{f(\mathbf{p})}{|\mathbf{p}|}$$

Equilibrium value

$$\frac{1}{g^2 \mu^2} \frac{dE}{dt} \approx 0.2$$

## Conclusions

- ▶ Energy loss is found as a solution of initial value problem
- ▶ Extremely prolate QGP is discussed as an example
- ▶ Strong time and directional dependence of  $dE/dx$  is demonstrated
- ▶  $dE/dx$  in unstable QGP is much bigger than in equilibrium one

for more see:

M. Carrington, K. Deja and St. Mrówczyński,  
Acta Physica Polonica B - Proceedings Supplement **5**, 047 (2012); *ibid* **5**, 343 (2012)