

Production of light nuclei in relativistic heavy-ion collisions

Stanisław Mrówczyński

*Institute of Physics, Jan Kochanowski University, Kielce, Poland
and National Centre for Nuclear Research, Warsaw, Poland*

Background


Conventional wisdom

- ▶ Production of light nuclei in relativistic heavy-ion collisions is well understood as a process of final state interactions.
- ▶ Coalescence models work well in a broad collision energy range.

News from RHIC & LHC

- ▶ Production of ^2H , $^2\bar{\text{H}}$, ^3H , $^3\bar{\text{H}}$, ^3He , $^3\bar{\text{He}}$, ^4He , $^4\bar{\text{He}}$, $_{\Lambda}^3\text{H}$, $_{\bar{\Lambda}}^3\bar{\text{H}}$ is observed in midrapidity.
- ▶ Matter-antimatter symmetry is seen.
- ▶ Thermal model properly describes yields of light nuclei.

Thermal model prediction



baryonless fireball

$$\text{Yield} \sim g e^{-\frac{m}{T}}$$

$$T = 156 \text{ MeV}$$


Can light nuclei exist in a fireball?

- ▶ Interparticle spacing in a hadron gas is about 1.5 fm at $T = 156$ MeV.
- ▶ Root mean square radius of a deuteron is 2.0 fm.
- ▶ Binding energy of a deuteron is 2.2 MeV.
- ▶ A hadron gas at $T = 156$ MeV is essentially a classical system.

- *Snowflakes in hell ?*
- *No, snowflakes from hell.*



Final state interaction – conventional approach to production of light nuclei




S.T. Butler & C.A. Pearson, Phys. Rev. **129**, 836 (1963)
A. Schwarzschild & C. Zupancic, Phys. Rev. **129**, 854 (1963)

Factorization of production of nucleons and formation of a nucleus

Deuteron production cross section

$$\frac{d\sigma^D}{d^3\mathbf{P}_D} = W \frac{d\sigma^{np}}{d^3\mathbf{p}_n d^3\mathbf{p}_p} \quad \frac{1}{2}\mathbf{P}_D = \mathbf{p}_n = \mathbf{p}_p$$

source - fireball



Deuteron formation rate

$$W = \frac{3}{4} (2\pi)^3 \int d^3\mathbf{r}_1 d^3\mathbf{r}_2 D(\mathbf{r}_1) D(\mathbf{r}_2) |\psi(\mathbf{r}_1, \mathbf{r}_2)|^2$$

spin factor

H. Sato and K. Yazaki, Phys. Lett. B 98 , 153 (1981)

Deuteron formation rate vs. n-p correlation

$$\psi(\mathbf{r}_1, \mathbf{r}_2) = e^{i\mathbf{P} \cdot \mathbf{R}} \varphi(\mathbf{r}) \quad \mathbf{R} \equiv \frac{1}{2}(\mathbf{r}_1 + \mathbf{r}_2), \quad \mathbf{r} \equiv \mathbf{r}_1 - \mathbf{r}_2$$

$$W = \frac{3}{4}(2\pi)^3 \int d^3\mathbf{r} D_r(\mathbf{r}) |\varphi(\mathbf{r})|^2$$

$$D_r(\mathbf{r}) \equiv \int d^3\mathbf{R} D\left(\mathbf{R} - \frac{1}{2}\mathbf{r}\right) D\left(\mathbf{R} + \frac{1}{2}\mathbf{r}\right)$$

n-p – correlation function

$$C(\mathbf{q}) = \int d^3\mathbf{r} D_r(\mathbf{r}) |\varphi_{\mathbf{q}}(\mathbf{r})|^2$$


$\varphi(\mathbf{r})$ – wave function of a bound state

$\varphi_{\mathbf{q}}(\mathbf{r})$ – wave function of a scattering state

St. Mrówczyński, Phys. Lett. B 277, 43 (1992)

Quantum-mechanical meaning of the formation rate formula

Sudden approximation



Transition matrix element

$$W = \left| \int d^3\mathbf{r} \psi^*(\mathbf{r}) \varphi(\mathbf{r}) \right|^2 = \int d^3\mathbf{r} d^3\mathbf{r}' \varphi^*(\mathbf{r}') \underbrace{\psi(\mathbf{r}') \psi^*(\mathbf{r}) \varphi(\mathbf{r})}_{\rho(\mathbf{r}', \mathbf{r})} \text{ density matrix}$$


$$W = \int d^3\mathbf{r} d^3\mathbf{r}' \varphi^*(\mathbf{r}') \rho(\mathbf{r}', \mathbf{r}) \varphi(\mathbf{r})$$

If density matrix is diagonal (random phase approximation)

$$\rho(\mathbf{r}', \mathbf{r}) = D(\mathbf{r}) \delta^{(3)}(\mathbf{r}' - \mathbf{r}) \quad \Rightarrow \quad W = \int d^3\mathbf{r} D(\mathbf{r}) |\varphi(\mathbf{r})|^2$$

Energy-momentum conservation

source - fireball



Nucleons are intermediate scattering states

$$E_i^2 - \mathbf{p}_i^2 \neq m_i^2 \quad i = n, p$$


Energy-momentum conservation

$$\left\{ \begin{array}{l} \mathbf{p}_p + \mathbf{p}_n = \mathbf{p}_D \\ E_p + E_n = E_D \end{array} \right.$$

St. Mrówczyński, J. Phys. G 11, 1087 (1987)

Thermal vs. coalescence model


- ▶ The two models usually give quantitatively similar predictions.
- ▶ How to falsify one of the models experimentally?



Karl Popper 1902-1994

The first idea: ${}^4\text{He}$ vs. ${}^4\text{Li}$

${}^4\text{He}$




$$r_{\text{RMS}} = 1.68 \text{ fm}$$

$$\varepsilon_B = 28.3 \text{ MeV}$$

$$m = 3727.4 \text{ MeV}$$

$$s = 0$$

${}^4\text{Li}$




$$\Gamma = 6 \text{ MeV}$$

$$m = m_{{}^3\text{He}} + m_p + 4.1 \text{ MeV}$$

$$m = 3749.7 \text{ MeV}$$

$$s = 2$$



Thermal model

$$\frac{\text{Yield}({}^4\text{Li})}{\text{Yield}({}^4\text{He})} = \frac{2S_{\text{Li}} + 1}{2S_{\text{He}} + 1} = 5$$




Coalescence model

$$\frac{\text{Yield}({}^4\text{Li})}{\text{Yield}({}^4\text{He})} = \frac{W_{\text{Li}}}{W_{\text{He}}}$$


Formation rates of ${}^4\text{He}$ & ${}^4\text{Li}$

$$W = g_s g_I (2\pi)^9 \int d^3\mathbf{r}_1 d^3\mathbf{r}_2 \int d^3\mathbf{r}_3 d^3\mathbf{r}_4 D(\mathbf{r}_1) D(\mathbf{r}_2) D(\mathbf{r}_3) D(\mathbf{r}_4) |\psi(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4)|^2$$



$$\mathbf{r}_{ij} \equiv \mathbf{r}_i - \mathbf{r}_j$$

$$|\psi_{\text{He}}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4)|^2 \sim \exp \left[-\alpha (\mathbf{r}_{12}^2 + \mathbf{r}_{13}^2 + \mathbf{r}_{14}^2 + \mathbf{r}_{23}^2 + \mathbf{r}_{24}^2 + \mathbf{r}_{34}^2) \right]$$



J. C. Bergstrom, Nucl. Phys. A 327, 458 (1979)

$$\mathbf{z} \equiv \mathbf{r}_4 - \frac{1}{3}(\mathbf{r}_1 + \mathbf{r}_2 + \mathbf{r}_3)$$

$$|\psi_{\text{Li}}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4)|^2 \sim \exp \left[-\beta (\mathbf{r}_{12}^2 + \mathbf{r}_{13}^2 + \mathbf{r}_{23}^2) \right] \mathbf{z}^4 \exp(-\gamma \mathbf{z}^2) |Y_{lm}(\Omega_z)|^2$$

Formation rates of ${}^4\text{He}$ & ${}^4\text{Li}$

$$W = g_s g_I (2\pi)^9 \int d^3\mathbf{r}_1 d^3\mathbf{r}_2 \int d^3\mathbf{r}_3 d^3\mathbf{r}_4 D(\mathbf{r}_1) D(\mathbf{r}_2) D(\mathbf{r}_3) D(\mathbf{r}_4) |\psi(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4)|^2$$

Source function

$$D(\mathbf{r}_i) = \frac{1}{(2\pi R_s^2)^{3/2}} \exp\left(-\frac{\mathbf{r}_i^2}{2R_s^2}\right) \quad i = 1, 2, 3, 4$$

If emission time included

Jacobi variables

$$\left\{ \begin{array}{l} \mathbf{R} \equiv \frac{1}{4}(\mathbf{r}_1 + \mathbf{r}_2 + \mathbf{r}_3 + \mathbf{r}_4) \\ \mathbf{x} \equiv \mathbf{r}_2 - \mathbf{r}_1 \\ \mathbf{y} \equiv \mathbf{r}_3 - \frac{1}{2}(\mathbf{r}_1 + \mathbf{r}_2) \\ \mathbf{z} \equiv \mathbf{r}_4 - \frac{1}{3}(\mathbf{r}_1 + \mathbf{r}_2 + \mathbf{r}_3) \end{array} \right. \quad R_s \rightarrow \sqrt{R_s^2 + v^2 \tau^2}$$

► $\mathbf{r}_1^2 + \mathbf{r}_2^2 + \mathbf{r}_3^2 + \mathbf{r}_4^2 = 4\mathbf{R}^2 + \frac{1}{2}\mathbf{x}^2 + \frac{2}{3}\mathbf{y}^2 + \frac{3}{4}\mathbf{z}^2$

► $\mathbf{r}_{12}^2 + \mathbf{r}_{13}^2 + \mathbf{r}_{14}^2 + \mathbf{r}_{23}^2 + \mathbf{r}_{24}^2 + \mathbf{r}_{34}^2 = 2\mathbf{x}^2 + \frac{8}{3}\mathbf{y}^2 + 3\mathbf{z}^2$

$$\mathbf{r}_{ij} \equiv \mathbf{r}_i - \mathbf{r}_j$$

Fully analytic calculations
are possible!

Formation rates of ${}^4\text{He}$ & ${}^4\text{Li}$

- ▶
$$W_{\text{He}} = \frac{\pi^{9/2}}{2^{9/2}} \frac{1}{(R_s^2 + R_\alpha^2)^{9/2}}$$
- ▶
$$W_{\text{Li}} = \frac{3\pi^{9/2}}{2^{11/2}} \left(\begin{array}{c} \frac{5}{2} \\ 1 \end{array} \right) \frac{R_s^4}{\left(R_s^2 + \frac{1}{2} R_c^2 \right)^3 \left(R_s^2 + \frac{4}{7} R_{\text{Li}}^2 - \frac{3}{7} R_c^2 \right)^{7/2}} \quad \left(\begin{array}{l} l=1 \\ l=2 \end{array} \right)$$

R_s – root mean square radius of the source


R_α – root mean square radius of ${}^4\text{He}$

R_{Li} – root mean square radius of ${}^4\text{Li}$

R_c – root mean square radius of ${}^3\text{He}$ cluster in ${}^4\text{Li}$


Ratio of yields of ^4Li to ^4He

In the thermal model the ratio equals 5.




How to observe ${}^4\text{Li}$?

Measurement of the correlation function of ${}^3\text{He}-p$ is needed



J. Pochodzala et al. Phys. Rev. C **35**, 1695 (1987)



T. A. Armstrong et al. Phys. Rev. C **65**, 014906 (2001)

The second idea: h - D correlations

- ▶ Hadron-deuteron correlations carry information about a source of deuterons.
- ▶ A measurement of K^- - D or p - D correlation functions can falsify the thermal or coalescence model.

Hadron-deuteron correlation function

1) Deuteron is treated as an elementary particle

Experimental definition

$$\frac{dN_{\pi D}}{d\mathbf{p}_\pi d\mathbf{p}_D} = R(\mathbf{p}_\pi, \mathbf{p}_D) \frac{dN_\pi}{d\mathbf{p}_\pi} \frac{dN_D}{d\mathbf{p}_D}$$

pion a generic hadron

Theoretical formula

$$R(\mathbf{p}_\pi, \mathbf{p}_D) = \int d^3r_\pi d^3r_D D(\mathbf{r}_\pi) D(\mathbf{r}_D) |\psi(\mathbf{r}_\pi, \mathbf{r}_D)|^2$$


distribution
of emission points π - D wave function

S.E. Koonin, Phys. Lett. B **70**, 43 (1977)

R. Lednicky and V.L. Lyuboshitz, Yad. Fiz. **35**, 1316 (1982)

Hadron-deuteron correlation function

1) Deuteron is treated as an elementary particle cont.

Center-of-mass variables

$$\left\{ \begin{array}{l} \mathbf{R} = \frac{m_\pi \mathbf{r}_\pi + m_D \mathbf{r}_D}{m_\pi + m_D} \\ \mathbf{r}_{\pi D} = \mathbf{r}_\pi - \mathbf{r}_D \end{array} \right. \quad \left\{ \begin{array}{l} \mathbf{P} = (\mathbf{p}_1 + \mathbf{p}_2) \\ \mathbf{q} = \frac{m_\pi \mathbf{p}_D - m_D \mathbf{p}_\pi}{m_\pi + m_D} \end{array} \right. \quad \psi(\mathbf{r}_\pi, \mathbf{r}_D) = e^{i \mathbf{P} \cdot \mathbf{R}} \phi_{\mathbf{q}}(\mathbf{r}_{\pi D})$$

$$R(\mathbf{q}) = \int d^3 r_{\pi D} D_r(\mathbf{r}_{\pi D}) |\phi_{\mathbf{q}}(\mathbf{r}_{\pi D})|^2$$

'Relative' source

$$D_r(\mathbf{r}_{\pi D}) \equiv \int d^3 \mathbf{R} D\left(\mathbf{R} - \frac{m_\pi}{m_\pi + m_D} \mathbf{r}_{\pi D}\right) D\left(\mathbf{R} + \frac{m_D}{m_\pi + m_D} \mathbf{r}_{\pi D}\right)$$

$$D(\mathbf{r}) = \left(\frac{1}{2\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{2R^2}\right) \quad \Rightarrow \quad D_r(\mathbf{r}) = \left(\frac{1}{4\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{4R^2}\right)$$

Hadron-deuteron correlation function

2) Deuteron is treated as a bound state of neutron and proton

Experimental definition

$$\frac{dN_{\pi D}}{d\mathbf{p}_\pi d\mathbf{p}_D} = R(\mathbf{p}_\pi, \mathbf{p}_D) A \frac{dN_\pi}{d\mathbf{p}_\pi} \frac{dN_n}{d\mathbf{p}_n} \frac{dN_p}{d\mathbf{p}_p}$$

$$\frac{1}{2}\mathbf{P}_D = \mathbf{p}_n = \mathbf{p}_p$$

Deuteron formation rate

$$\frac{dN_D}{d\mathbf{p}_D} = A \frac{dN_n}{d\mathbf{p}_n} \frac{dN_p}{d\mathbf{p}_p}$$

$$A = \frac{3}{4} (2\pi)^3 \int d^3\mathbf{r}_n d^3\mathbf{r}_p D(\mathbf{r}_n) D(\mathbf{r}_p) |\psi_D(\mathbf{r}_n, \mathbf{r}_p)|^2 = \frac{3}{4} (2\pi)^3 \int d^3r_{np} D_r(\mathbf{r}_{np}) |\phi_D(\mathbf{r}_{np})|^2$$

↑
spin factor
↑
 $\psi_D(\mathbf{r}_n, \mathbf{r}_p) = e^{i\mathbf{P}\cdot\mathbf{R}} \phi_D(\mathbf{r}_{np})$

Hadron-deuteron correlation function

2) Deuteron is treated as a bound state of neutron and proton cont.

Theoretical formula

$$R(\mathbf{p}_\pi, \mathbf{p}_D) = \frac{1}{A} \int d^3 r_\pi d^3 r_n d^3 r_p D(\mathbf{r}_\pi) D(\mathbf{r}_n) D(\mathbf{r}_p) |\psi_{\pi D}(\mathbf{r}_\pi, \mathbf{r}_n, \mathbf{r}_p)|^2$$

Center-of-mass (Jacobi) variables

$$\left\{ \begin{array}{l} \mathbf{R} = \frac{m_\pi \mathbf{r}_\pi + m_n \mathbf{r}_n + m_p \mathbf{r}_p}{m_\pi + m_n + m_p} \\ \mathbf{r}_{np} = \mathbf{r}_n - \mathbf{r}_p \\ \mathbf{r}_{\pi D} = \mathbf{r}_\pi - \frac{m_n \mathbf{r}_n + m_p \mathbf{r}_p}{m_n + m_p} \end{array} \right.$$

$$\psi_{\pi D}(\mathbf{r}_\pi, \mathbf{r}_n, \mathbf{r}_p) = e^{i \mathbf{P} \cdot \mathbf{R}} \phi_{\mathbf{q}}(\mathbf{r}_{\pi D}) \phi_D(\mathbf{r}_{np})$$

$$D_{3r}(\mathbf{r}_{\pi D}) D_r(\mathbf{r}_{np}) \equiv \int d^3 \mathbf{R} D(\mathbf{r}_\pi) D(\mathbf{r}_n) D(\mathbf{r}_p)$$

$$D(\mathbf{r}) = \left(\frac{1}{2\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{2R^2}\right)$$

Deuteron formation rate cancels out!

$$R(\mathbf{q}) = \int d^3 r_{\pi D} D_{3r}(\mathbf{r}_{\pi D}) |\phi_{\mathbf{q}}(\mathbf{r}_{\pi D})|^2$$

$$D_r(\mathbf{r}) = \left(\frac{1}{4\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{4R^2}\right)$$

$$D_{3r}(\mathbf{r}) = \left(\frac{1}{3\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{3R^2}\right)$$


Thermal vs. coalescence model



Thermal model

$$R(\mathbf{q}) = \int d^3 r_{\pi D} D_r(\mathbf{r}_{\pi D}) |\phi_{\mathbf{q}}(\mathbf{r}_{\pi D})|^2$$

$$D(\mathbf{r}) = \left(\frac{1}{2\pi R^2} \right)^{3/2} \exp \left(-\frac{\mathbf{r}^2}{2R^2} \right)$$



$$D_r(\mathbf{r}) = \left(\frac{1}{4\pi R^2} \right)^{3/2} \exp \left(-\frac{\mathbf{r}^2}{4R^2} \right)$$

$$D_{3r}(\mathbf{r}) = \left(\frac{1}{3\pi R^2} \right)^{3/2} \exp \left(-\frac{\mathbf{r}^2}{3R^2} \right)$$



Coalescence model

$$R(\mathbf{q}) = \int d^3 r_{\pi D} D_{3r}(\mathbf{r}_{\pi D}) |\phi_{\mathbf{q}}(\mathbf{r}_{\pi D})|^2$$



Computation of correlation function

$$R(\mathbf{q}) = \int d^3r_{\pi D} D_r(\mathbf{r}_{\pi D}) |\phi_{\mathbf{q}}(\mathbf{r}_{\pi D})|^2$$

Source function

$$D_r(\mathbf{r}) = \left(\frac{1}{4\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{4R^2}\right) \quad \text{or} \quad D_{3r}(\mathbf{r}) = \left(\frac{1}{3\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{3R^2}\right)$$

Wave function

$$\phi_{\mathbf{q}}(\mathbf{r}) = e^{i\mathbf{qr}} + f(q) \frac{e^{iqr}}{r}$$

S-wave amplitude

$$f(q) = -\frac{a}{1 + iqa}$$

a - scattering length

Coulomb effect


$$R(\mathbf{q}) \rightarrow G(\mathbf{q})R(\mathbf{q})$$

Gamov factor

$$G(\mathbf{q}) = \pm \frac{2\pi}{a_B q} \frac{1}{\exp\left(\frac{2\pi}{a_B q}\right) - 1}$$

a_B - Bohr radius


K^- - D correlation functions



$$a = (1.46 - 1.08i) \text{ fm}$$

$$2.00 = \sqrt{\frac{4}{3}} 1.73 = \frac{4}{3} 1.50$$

***p-D* correlation functions**



$$a_{1/2} = 4.0 \text{ fm}$$
$$a_{3/2} = 11.0 \text{ fm}$$

$$R(q) = \frac{1}{3} R_{1/2}(q) + \frac{2}{3} R_{3/2}(q)$$

$$2.00 = \sqrt{\frac{4}{3}} 1.73 = \frac{4}{3} 1.50$$

Conclusions

${}^4\text{He}$ vs. ${}^4\text{Li}$

- ▶ The thermal and coalescence models give different predictions on the ratio of yields of ${}^4\text{Li}$ to ${}^4\text{He}$.
- ▶ In the thermal model the ratio of yields is independent of collision centrality.
- ▶ In the coalescence model the ratio is maximal for central collisions and rapidly decreases when one goes to peripheral collisions.
- ▶ Since ${}^4\text{Li}$ can be observed through the correlation function of ${}^3\text{He}-p$, the correlation needs to be computed.

h - D correlations

- ▶ Hadron-deuteron correlations carry information about source of deuterons.
- ▶ Measurement of h - D correlation function can tell us whether deuterons are directly emitted from a fireball or deuterons are formed due to final state interactions.
- ▶ K^- - D & p - D correlation functions show a sufficient sensitivity to a size of particle source to falsify the thermal or coalescence model.