

Hadron-Deuteron Correlations & Production of Light Nuclei in Relativistic Heavy-Ion Collisions


Stanisław Mrówczyński

*Institute of Physics, Jan Kochanowski University, Kielce, Poland
and National Centre for Nuclear Research, Warsaw, Poland*

In collaboration with **Patrycja Słoń**

Background

- ▶ Production of ${}^2\text{H}$, ${}^2\bar{\text{H}}$, ${}^3\text{H}$, ${}^3\bar{\text{H}}$, ${}^3\text{He}$, ${}^3\bar{\text{He}}$, ${}^4\text{He}$, ${}^4\bar{\text{He}}$, ${}^3_{\Lambda}\text{H}$, ${}^3_{\Lambda}\bar{\text{H}}$ is observed in midrapidity at RHIC & LHC.
- ▶ Thermal model properly describes yields of light nuclei.



baryonless fireball

$$\text{Yield} \sim g e^{-\frac{m}{T}}$$

$$T = 156 \text{ MeV}$$

A. Andronic, P. Braun-Munzinger, K. Redlich and J. Stachel, *Nature* **561**, 321 (2018)


Can light nuclei exist in a fireball?

- ▶ Interparticle spacing in a hadron gas is about 1.5 fm at $T = 156$ MeV.
- ▶ Root mean square radius of a deuteron is 2.0 fm.
- ▶ Binding energy of a deuteron is 2.2 MeV.
- ▶ A hadron gas at $T = 156$ MeV is essentially a classical system.

- Snowflakes in hell ?
- No, snowflakes from hell.




Final state interaction – conventional approach to production of light nuclei



S.T. Butler & C.A. Pearson, Phys. Rev. **129**, 836 (1963)
A. Schwarzschild & C. Zupancic, Phys. Rev. **129**, 854 (1963)
H. Sato and K. Yazaki, Phys. Lett. B **98**, 153 (1981)

Thermal vs. coalescence model


- ▶ The two models usually give quantitatively similar predictions.
- ▶ How to falsify one of the models experimentally?



Karl Popper 1902-1994


${}^4\text{He}$ vs. ${}^4\text{Li}$

${}^4\text{He}$



$r_{\text{RMS}} = 1.68 \text{ fm}$
 $\varepsilon_B = 28.3 \text{ MeV}$
 $m = 3727.4 \text{ MeV}$
 $s = 0$

${}^4\text{Li}$



${}^4\text{Li} \rightarrow {}^3\text{He} + p$
 $\Gamma = 6 \text{ MeV}$
 $m = m_{{}^3\text{He}} + m_p + 4.1 \text{ MeV}$
 $m = 3749.7 \text{ MeV}$
 $s = 2$

► Thermal model $\frac{\text{Yield}({}^4\text{Li})}{\text{Yield}({}^4\text{He})} = \frac{2S_{\text{Li}} + 1}{2S_{\text{He}} + 1} = 5$

► Coalescence model $\frac{\text{Yield}({}^4\text{Li})}{\text{Yield}({}^4\text{He})} \sim 1$ and strongly centrality dependent

Sylwia's talk on Sunday

The second idea

- ▶ Hadron-deuteron correlations carry information about a source of deuterons.
- ▶ A measurement of K^- - D or p - D correlation functions is suggested to falsify the thermal or coalescence model.

Hadron-deuteron correlation function

1) Deuteron is treated as an elementary particle

Experimental definition

$$\frac{dN_{\pi D}}{d\mathbf{p}_{\pi} d\mathbf{p}_D} = R(\mathbf{p}_{\pi}, \mathbf{p}_D) \frac{dN_{\pi}}{d\mathbf{p}_{\pi}} \frac{dN_D}{d\mathbf{p}_D}$$

pion a generic hadron

Theoretical formula

$$R(\mathbf{p}_{\pi}, \mathbf{p}_D) = \int d^3 r_{\pi} d^3 r_D D(\mathbf{r}_{\pi}) D(\mathbf{r}_D) |\psi(\mathbf{r}_{\pi}, \mathbf{r}_D)|^2$$

distribution
of emission points

π -D wave function

S.E. Koonin, Phys. Lett. B **70**, 43 (1977)

R. Lednicky and V.L. Lyuboshitz, Yad. Fiz. **35**, 1316 (1982)

Hadron-deuteron correlation function

1) Deuteron is treated as an elementary particle cont.

Center-of-mass variables

$$\left\{ \begin{array}{l} \mathbf{R} = \frac{m_\pi \mathbf{r}_\pi + m_D \mathbf{r}_D}{m_\pi + m_D} \\ \mathbf{r}_{\pi D} = \mathbf{r}_\pi - \mathbf{r}_D \end{array} \right. \quad \left\{ \begin{array}{l} \mathbf{P} = (\mathbf{p}_1 + \mathbf{p}_2) \\ \mathbf{q} = \frac{m_\pi \mathbf{p}_D - m_D \mathbf{p}_\pi}{m_\pi + m_D} \end{array} \right. \quad \psi(\mathbf{r}_\pi, \mathbf{r}_D) = e^{i\mathbf{P}\mathbf{R}} \phi_{\mathbf{q}}(\mathbf{r}_{\pi D})$$

$$R(\mathbf{q}) = \int d^3 r_{\pi D} D_r(\mathbf{r}_{\pi D}) \left| \phi_{\mathbf{q}}(\mathbf{r}_{\pi D}) \right|^2$$

'Relative' source

$$D_r(\mathbf{r}_{\pi D}) \equiv \int d^3 \mathbf{R} D\left(\mathbf{R} - \frac{m_\pi}{m_\pi + m_D} \mathbf{r}_{\pi D}\right) D\left(\mathbf{R} + \frac{m_D}{m_\pi + m_D} \mathbf{r}_{\pi D}\right)$$

$$D(\mathbf{r}) = \left(\frac{1}{2\pi R^2}\right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{2R^2}\right) \quad \Rightarrow \quad D_r(\mathbf{r}) = \left(\frac{1}{4\pi R^2}\right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{4R^2}\right)$$

Hadron-deuteron correlation function

2) Deuteron is treated as a bound state of neutron and proton

Experimental definition

$$\frac{dN_{\pi D}}{d\mathbf{p}_\pi d\mathbf{p}_D} = R(\mathbf{p}_\pi, \mathbf{p}_D) A \frac{dN_\pi}{d\mathbf{p}_\pi} \frac{dN_n}{d\mathbf{p}_n} \frac{dN_p}{d\mathbf{p}_p}$$

$$\frac{1}{2} \mathbf{P}_D = \mathbf{p}_n = \mathbf{p}_p$$

Deuteron formation rate

$$\frac{dN_D}{d\mathbf{p}_D} = A \frac{dN_n}{d\mathbf{p}_n} \frac{dN_p}{d\mathbf{p}_p}$$

$$A = \frac{3}{4} (2\pi)^3 \int d^3\mathbf{r}_n d^3\mathbf{r}_p D(\mathbf{r}_n) D(\mathbf{r}_p) |\psi_D(\mathbf{r}_n, \mathbf{r}_p)|^2 = \frac{3}{4} (2\pi)^3 \int d^3r_{np} D_r(\mathbf{r}_{np}) |\phi_D(\mathbf{r}_{np})|^2$$

spin factor

$$\psi_D(\mathbf{r}_n, \mathbf{r}_p) = e^{i\mathbf{P}\mathbf{R}} \phi_D(\mathbf{r}_{np})$$

Hadron-deuteron correlation function

2) Deuteron is treated as a bound state of neutron and proton cont.

Theoretical formula

$$R(\mathbf{p}_\pi, \mathbf{p}_D) = \frac{1}{A} \int d^3 r_\pi d^3 r_n d^3 r_p D(\mathbf{r}_\pi) D(\mathbf{r}_n) D(\mathbf{r}_p) \left| \psi_{\pi D}(\mathbf{r}_\pi, \mathbf{r}_n, \mathbf{r}_p) \right|^2$$

Center-of-mass (Jacobi) variables

$$\left\{ \begin{array}{l} \mathbf{R} = \frac{m_\pi \mathbf{r}_\pi + m_n \mathbf{r}_n + m_p \mathbf{r}_p}{m_\pi + m_n + m_p} \\ \mathbf{r}_{np} = \mathbf{r}_n - \mathbf{r}_p \\ \mathbf{r}_{\pi D} = \mathbf{r}_\pi - \frac{m_n \mathbf{r}_n + m_p \mathbf{r}_p}{m_n + m_p} \end{array} \right.$$

$$\psi_{\pi D}(\mathbf{r}_\pi, \mathbf{r}_n, \mathbf{r}_p) = e^{i\mathbf{P}\mathbf{R}} \phi_q(\mathbf{r}_{\pi D}) \phi_D(\mathbf{r}_{np})$$

$$D_{3r}(\mathbf{r}_{\pi D}) D_r(\mathbf{r}_{np}) \equiv \int d^3 \mathbf{R} D(\mathbf{r}_\pi) D(\mathbf{r}_n) D(\mathbf{r}_p)$$

$$D(\mathbf{r}) = \left(\frac{1}{2\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{2R^2} \right)$$

Deuteron formation rate cancels out!

$$R(\mathbf{q}) = \int d^3 r_{\pi D} D_{3r}(\mathbf{r}_{\pi D}) \left| \phi_q(\mathbf{r}_{\pi D}) \right|^2$$

$$D_r(\mathbf{r}) = \left(\frac{1}{4\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{4R^2} \right)$$



$$D_{3r}(\mathbf{r}) = \left(\frac{1}{3\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{3R^2} \right)$$

Thermal vs. coalescence model

► Thermal model

$$R(\mathbf{q}) = \int d^3 r_{\pi D} D_r(\mathbf{r}_{\pi D}) \left| \phi_{\mathbf{q}}(\mathbf{r}_{\pi D}) \right|^2$$

$$D(\mathbf{r}) = \left(\frac{1}{2\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{2R^2} \right)$$



$$D_r(\mathbf{r}) = \left(\frac{1}{4\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{4R^2} \right)$$

$$D_{3r}(\mathbf{r}) = \left(\frac{1}{3\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{3R^2} \right)$$



► Coalescence model

$$R(\mathbf{q}) = \int d^3 r_{\pi D} D_{3r}(\mathbf{r}_{\pi D}) \left| \phi_{\mathbf{q}}(\mathbf{r}_{\pi D}) \right|^2$$

Computation of correlation function

$$R(\mathbf{q}) = \int d^3 r_{\pi D} D_r(\mathbf{r}_{\pi D}) \left| \phi_{\mathbf{q}}(\mathbf{r}_{\pi D}) \right|^2$$

Source function

$$D_r(\mathbf{r}) = \left(\frac{1}{4\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{4R^2} \right) \quad \text{or} \quad D_{3r}(\mathbf{r}) = \left(\frac{1}{3\pi R^2} \right)^{3/2} \exp\left(-\frac{\mathbf{r}^2}{3R^2} \right)$$

Wave function

$$\phi_{\mathbf{q}}(\mathbf{r}) = e^{i\mathbf{q}\mathbf{r}} + f(q) \frac{e^{iqr}}{r}$$

S-wave amplitude

$$f(q) = -\frac{a}{1 + iqa}$$

a - scattering length

Coulomb effect


$$R(\mathbf{q}) \rightarrow G(\mathbf{q})R(\mathbf{q})$$

Gamov factor

$$G(\mathbf{q}) = \pm \frac{2\pi}{a_B q} \frac{1}{\exp\left(\frac{2\pi}{a_B q} \right) - 1}$$


a_B - Bohr radius

$K^- - D$ correlation functions



$$a = (1.46 - 1.08i) \text{ fm} \quad 2.00 = \sqrt{\frac{4}{3}} 1.73 = \frac{4}{3} 1.50$$

p-D correlation functions



$$a_{1/2} = 4.0 \text{ fm}$$

$$a_{3/2} = 11.0 \text{ fm}$$

$$R(q) = \frac{1}{3} R_{1/2}(q) + \frac{2}{3} R_{3/2}(q)$$

$$2.00 = \sqrt{\frac{4}{3}} 1.73 = \frac{4}{3} 1.50$$

Conclusions

- ▶ Hadron-deuteron correlations carry information about source of deuterons.
- ▶ Measurement of h - D correlation function can tell us whether deuterons are directly emitted from a fireball or deuterons are formed due to final state interactions.
- ▶ K^- - D & p - D correlation functions show a sufficient sensitivity to a size of particle source to falsify the thermal or coalescence model.